


ARJUNA 2.0 (JEE)

Motion in a Straight Line


DPP-12

1. The velocity (v) of a particle varies with position (x) as shown in the graph. Its acceleration when $x = 4$ m will be




- (A) -8 m/s^2 (B) 8 m/s^2
(C) 2 m/s^2 (D) -2 m/s^2

2. From the velocity-time graph of a particle moving in a straight line. The ratio of average velocity for interval 3 s and instantaneous velocity at 3 s.




- (A) 1 : 1 (B) 1 : 2
(C) 7 : 3 (D) 3 : 2

3. The velocity (v) versus time (t) graph of a particle moving along a straight line is as shown in the given figure. During time interval $t = 0$ to $t = 6$ s, the magnitude of displacement and the distance travelled are in the ratio of




- (A) 1 : 1 (B) 1 : 3
(C) 2 : 3 (D) 3 : 1

4. The position-time graph of two particles P and Q are as shown in figure. The ratio of their velocities $\frac{V_P}{V_Q}$ is




- (A) 1 : 3 (B) $\sqrt{3} : 1$
(C) 3 : 1 (D) $1 : \sqrt{3}$

5. The figure shows position (x) versus time (t) graph of a particle moving along x -axis. During time interval $t = 0$ to $t = 4$ s, how many times the particle comes to rest?




- (A) 4 (B) 2
(C) 3 (D) 5

6. Displacement-time ($x-t$) graph of a particle moving along a straight line path is shown in figure. Average speed of particle in the time interval 0 to 5 second is




- (A) 2 m/s (B) 12 m/s
(C) 10 m/s (D) 14 m/s

7. A particle is moving along positive x -axis with some initial velocity. The acceleration time graphs are shown. In which case velocity of particle will increased for entire time between t_1 to t_2 ?




- (A) Only in II (B) In I and III
(C) In I and II (D) In I, II and III

8. The velocity versus time graph of a body is shown in figure. If the slope of line is K then distance travelled by the body in time T is




- (A) $\frac{K^2 T}{2}$ (B) $\frac{KT^2}{2}$
(C) $\frac{K}{2T}$ (D) $\frac{T}{2K}$

9. A particle moves with an initial velocity $v_0 = 5\text{ m/s}$ in a straight line. If its acceleration a varies with time t as shown in a - t graph, then velocity of the particle at $t = 7\text{ s}$ is




- (A) 10 m/s (B) -10 m/s
(C) -14 m/s (D) 24 m/s

10. A particle starts from rest and is acted upon by a variable acceleration as shown. Velocity of the particle is maximum at time t is equal to




- (A) 5 s (B) 10 s
(C) 12 s (D) 18 s

11. Acceleration (a) versus time (t) graph of a particle moving along x -axis is as shown in the figure. Change in velocity of the body in the interval 0 to 15 s is



- (A) 20 m/s (B) 15 m/s
(C) 5 m/s (D) 10 m/s

12. Acceleration (a) versus time (t) graph of a particle started from rest is as shown in figure




Which point velocity of the particle is maximum?

- (A) A (B) B
(C) C (D) All of these

ANSWERS KEY

1. (A)
2. (B)
3. (B)
4. (A)
5. (A)
6. (D)
7. (D)
8. (B)
9. (C)
10. (C)
11. (D)
12. (C)



Any issue with DPP, please report by clicking here- <https://forms.gle/t2SzQVvQcs638c4r5>

Please share your feedback on PW Teachers- <https://forms.gle/jEBFswBuki4Ut2Lk6>



PW Mobile APP : <https://physicswala.page.link/?type=contact-us&data=open>

For PW Website : <https://www.physicswallah.live/contact-us>